

High-Resolution K' Imaging of the $z = 1.786$ Radio Galaxy 3C 294

Alan Stockton¹, Gabriela Canalizo¹

Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822

Susan E. Ridgway

Department of Physics and Astronomy, Johns Hopkins University, Homewood Campus,
Baltimore, MD 21218

ABSTRACT

We have obtained imaging in the K' band ($\sim I$ -band rest frame) of the $z = 1.786$ radio galaxy 3C 294 with the 36-element curvature-sensing adaptive optics system *Hokupa'a* and the Canada-France-Hawaii Telescope. At a resolution of $\lesssim 0''.15$, the galaxy is seen as a group of small but resolved knots distributed over a roughly triangular region $\sim 1''.4$ across. The interpretation of the structure depends on the location of the nucleus, as indicated by the compact radio core. Its position is uncertain by $\gtrsim 0''.5$ (2σ) because of uncertainties in the optical astrometry, but our best estimate places it at or near the southern apex of the distribution. If this location is correct, the most likely interpretation is that of a hidden quasar nucleus illuminating dusty infalling dwarf-galaxy-like clumps having characteristic sizes of ~ 1.5 kpc.

Subject headings: galaxies: active — galaxies: evolution — galaxies: formation
— galaxies: individual (3C 294) — radio continuum: galaxies

1. Introduction

High-redshift radio galaxies probably mark out regions of higher than average density in the early Universe, and they give us a window into formation processes for at least one kind of massive galaxy. Recent *Hubble Space Telescope* (*HST*) imaging of radio galaxies with redshifts > 2 in the rest-frame ultraviolet show that most comprise several components

¹Visiting Astronomer, Canada-France-Hawaii Telescope, operated by the National Research Council of Canada, the Centre National de la Recherche Scientifique de France, and the University of Hawaii.

and that the ultraviolet flux from these components is apparently dominated by recent star formation (van Breugel *et al.* 1998; Pentericci *et al.* 1998, 1999).

3C 294, at $z = 1.786$, is one of the dozen radio galaxies in the 3CR catalog with $z > 1.5$, so it is one of the most powerful radio sources in the observed Universe as well as a likely example of a massive galaxy in its youth. It has strong Ly- α emission extending over $\sim 10''$, roughly aligned with the radio structure (McCarthy *et al.* 1990). It also has a $V = 12$ star $< 10''$ to the west (Kristian, Sandage, & Katem 1974), which initially hampered optical and IR observations, but which now can be put to good use as an adaptive optics (AO) reference. In this *Letter* we describe the results of an initial AO imaging investigation of 3C 294.

2. Observations and Data Reduction

The AO observations were obtained with the University of Hawaii AO system *Hokupa‘a* (Graves *et al.* 1998) mounted at the f/36 focus of the Canada-France-Hawaii Telescope (CFHT). This system uses the curvature-sensing approach pioneered by François Roddier (Roddier, Northcott, & Graves 1991). Briefly, an image of the telescope primary is formed on a 36-element deformable mirror. Light from this mirror shortwards of $\sim 1 \mu\text{m}$ is sent by a beamsplitter to a membrane mirror, which is driven at 2.6 kHz to image extrafocal images on both sides of focus onto a 36-element avalanche-photodiode array. Corrections for the wavefront errors derived from the difference of these extrafocal images are sent back to the deformable bimorph mirror, which is updated at 1.3 kHz. Under typical seeing conditions at CFHT, and for sufficiently bright ($R \sim 12$) stars, diffraction-limited imaging can be achieved as short as I band (Graves *et al.* 1998). Our imaging of 3C 294 was in the K' band (Wainscoat & Cowie 1992), so the correction was excellent and quite stable over the course of the observations. The detector system was the University of Hawaii Quick Infrared Camera (QUIRC), which uses a 1024×1024 HAWAII array (Hodapp *et al.* 1996). We obtained 8 300 s exposures on 1998 July 3 UT and 14 300 s exposures on 1998 July 4 UT. Unusually rapid variation of the airglow emission compromised the reduction of data from the first night, and we use here only the data from the second night. The individual exposures were dithered in a pattern that kept the bright guide star off the edge of the detector for all exposures. The images were reduced using our standard iterative procedure (Stockton, Canalizo, & Close 1998). Briefly, we make a bad-pixel mask from a combination of hot pixels (from dark frames) and low-response pixels (from flat-field frames); these pixels are excluded from all sums and averages. For each dark-subtracted frame, 3 or 4 time-adjacent dithered frames were median averaged to make a sky frame,

which was normalized to the sky value of the frame in question and subtracted; then the residual was divided by a flat-field frame. These first-pass images were registered to the nearest pixel and median averaged. After a slight smoothing, this rough combined frame was used to generate object masks for each frame, which were then combined with the bad-pixel mask. This process was repeated to give better sky frames, better offsets for alignment, and a new combined image. This new image was used to replace bad pixels in each flattened frame with the median from the other frames, so that bad pixels would not affect the centering algorithm used to calculate the offsets. The final combined image was a straight average of the corrected, subpixel-registered frames, using a sigma-clipping algorithm in addition to the bad-pixel mask to eliminate deviant pixel values. Normally, we use field stars for registration, but in this case we had to use the ghost image of the guide star, since there were no other objects visible on individual frames. This ghost image was produced by a secondary reflection in the AO system optics; it was about 10 mag fainter than the guide star, and it was positioned $0''.114$ south and $7''.85$ east of the star. The image scale with the K' filter was determined to be $0''.03537 \pm 0''.00005 \text{ pixel}^{-1}$, based on an accurate measurement of the scale in the J filter by Claude Roddier and a determination of the ratio of the K' to J scales for the QUIRC camera from contemporaneous imaging data obtained with the University of Hawaii 2.2 m telescope.

3. Results

Our K' band image of 3C 294 samples the rest-frame spectrum from 6850–8300 Å, a region unlikely to be dominated by emission lines (the strongest expected lines in this bandpass are [Ar III] $\lambda\lambda 7136, 7751$ and [O II] $\lambda\lambda 7320, 7330$). This region is also close to the peak of the spectral-energy distribution (SED) for a stellar population with an age of $\gtrsim 2$ Gyr. If the luminosity at the center of this galaxy is dominated by a central bulge of relatively old stars, we should be able to see it. What we actually do see is shown in Fig. 1. Scattered light and a diffraction spike from the guide star extend from the lower right across the lower middle of the frame, and the ghost image of the guide star lies just above the diffraction spike. 3C 294 is at the middle of the frame. The structure is knotty and filamentary, comprising several distinct components within a roughly triangular region about $1''.4$ (~ 10 kpc, for $H_0 = 75$, $q_0 = 0.3$, $\Lambda = 0$, which we assume throughout this *Letter*) across, no one of which is clearly dominant. The data from the first night, although of lower quality, confirm the main features seen in Fig. 1.

Since, at larger scales, both the inner radio structure and the extended Ly- α emission are aligned along $\text{PA} \sim 20^\circ$ (McCarthy *et al.* 1990), one possible interpretation is that we

are seeing a dust scattering nebula centered on this same axis (Chambers 1999*a,b*). If this is the case, the hidden nucleus must lie somewhere near the faint southern tip of the “triangle”. However, it is also possible that we are seeing an assemblage of small merging components, and that the active nucleus is coincident with one of the peaks in our image. Thus, the interpretation of the observed structure is critically dependent on the location of the nucleus.

McCarthy *et al.* (1990) found a flat-spectrum central compact radio component, presumably to be identified with the nucleus, and they determined its position to a precision of ~ 50 mas. Because we have determined the position of ghost image and the image scale quite accurately, we can relate any point in our field to the position of the guide star to a similar precision. However, in order to relate our image to the radio position, we need to have high-quality astrometric positions for both the guide star and the radio nucleus in the same reference frame. There are two problems with accomplishing this task: (1) the McCarthy *et al.* (1990) radio data is referenced to the equinox B1950, epoch 1979.9 VLA calibrator reference frame, and converting to the equinox J2000, epoch 2000.0 FK5 frame is not completely straightforward; and (2) as McCarthy *et al.* (1990) point out, the positions for our guide star given by Veron (1966), Kristian *et al.* (1974), and Riley *et al.* (1980) do not agree very well. Our separate AO imaging of the guide star itself shows an added minor complication: the star is a double with a separation of $0''.13$ (see inset in Fig. 1).

We have converted the position of the central radio component to the FK5 frame by an empirical mapping of the B1950 VLA calibrator reference frame in the region of the source to the J2000 VLA calibrator frame. We first do a standard precession of B1950 calibrator positions from equinox B1950 to nominal equinox J2000, for calibrators within 15° of 3C 294. We then take the difference between these precessed positions and the cataloged J2000 VLA calibrator positions, and we fit a low-order reference surface to these differences (which were always $0''.5$ or less). The values of the corrections at the position of 3C 294 are then applied to the precessed B1950 VLA position for the central compact radio source. We obtain a position $14^{\text{h}}06^{\text{m}}44^{\text{s}}.077 \pm 0^{\text{s}}.002$, $+34^\circ11'24''.956 \pm 0''.025$ (J2000), where the quoted errors are the standard errors of the fit of the correction surface to the coordinate residuals. In addition, there is an uncertainty due to errors in the J2000 VLA calibrator positions, which is likely to be $\sim 0''.1$ or less.

Unfortunately, the AO guide star is too faint to appear in the Hipparcos/Tycho databases. In order to obtain a more accurate position for it, we have obtained large-image-scale, short-exposure CCD frames covering an $8'$ field including the star and have derived a plate solution from 12 USNO-A2.0 stars (after deleting 3 additional stars with outlying residuals). The residuals for the solution give a standard error of $0''.24$ in RA and $0''.34$ in

declination, so the 2σ uncertainty in the position of an individual star is over $0''.5$. For the mean position of the components of the AO guide star, we obtain $14^{\text{h}}06^{\text{m}}43^{\text{s}}.35, +34^{\circ}11'24''.2$ (J2000).

The USNO-A2.0 catalog is in the FK5 system tied to the Hipparcos reference frame. With the AO guide star and 3C 294 radio core positions, and our knowledge of the position of the AO guide star relative to its ghost image, we can locate the position of the radio core on our image. This position, with 2σ error bars (determined mostly by the uncertainty of the position of the guide star), is shown in the lower-left inset in Fig. 1, along with radio-core positions based on positions for the guide star given previously by Veron (1966), Kristian *et al.* (1974), and Riley *et al.* (1980). The position we derive for the radio core is consistent with a location at or near the southern tip of the K' structure, and apparently inconsistent with association with any of the three brightest blobs along the northern edge of the structure.

In a $3''$ -diameter synthetic aperture, we obtain a total K' magnitude for 3C 294 of 18.3 ± 0.3 . Lower-resolution imaging at the NASA Infrared Telescope Facility gives $K' = 18.3 \pm 0.1$. Both of these measurements are consistent with the $K = 18.0 \pm 0.3$ found by McCarthy *et al.* (1990) for a similar aperture.

4. Discussion

Radio galaxies at high redshifts often show multiple components, and these are often found to be aligned with the radio structure (*e.g.*, Pentericci *et al.* 1999, and references therein). This conclusion is based mostly on optical data, which corresponds to rest-frame UV for these objects, although recently some deep ground-based IR imaging has been presented by van Breugel *et al.* (1998), and programs involving HST NICMOS imaging of high-redshift radio galaxies are in progress (*e.g.*, Fosbury *et al.* 1999). One reason that 3C 294 is especially of interest in this context is that its redshift places the K' band in a region of the spectrum that is unlikely to be dominated by emission-line radiation or nebular thermal continuum. Since there is also no evidence in the VLA map of McCarthy *et al.* (1990) of radio counterparts to the rest-frame optical structure we see, the emission is most likely due either to stars or to scattering from a hidden quasar. The position we obtain for the flat-spectrum radio component of 3C 294 favors a location at or near the southern apex of the observed distribution of bright knots, and it therefore supports (though it cannot prove) the interpretation of the observed structure as an illumination cone, most likely due to dust scattering of radiation from a quasar nucleus. Figure 1 can be compared with Fig. 2 of McCarthy *et al.* (1990), which shows contours of the Ly- α and radio emission

(the short arrows in our Fig. 1 are at the positions of the radio knots K_N and K_S shown in their figure). The extended Ly- α emission also shows a well-defined triangular structure extending to the north, which McCarthy *et al.* (1990) suggest is due to anisotropic emission of ionizing radiation from a central non-thermal source. While the inferred Ly- α and K' cones are fairly well aligned, the Ly- α material extends $\sim 4''$, or roughly 4 times as far as does the continuum emission in our K' AO image. The Ly- α emission also extends to the other side of the radio nucleus, in a weak and somewhat poorly-defined “counter cone.” We do not see a corresponding feature at K' , but our dynamic range is not sufficient to put very strong limits on the presence of such material. If the illumination is truly biconical and intrinsically fairly symmetric, the southern cone must suffer significant extinction along our line of sight.

However, granting that the morphology we see in our AO image is likely to be at least partly determined by illumination effects, the material being illuminated also does seem to have an intrinsic distribution in small ($\sim 0''.2 \approx 1.5$ kpc) coherent bodies, which (on the not unreasonable assumption that they contain stars as well as dust) must shortly merge together. In fact, the tendency of these objects to show elongation roughly aligned in the direction of the apex may well be due to tidal stretching. Thus, the alternatives of illumination effects and merging subunits need not be starkly opposed to each other in this case, although the presence of an illumination cone, if confirmed (say, by polarization measurements), means that we are likely seeing only part of the action.

In summary, 3C 294 appears to be a particularly good example of several aspects of an emerging picture of high-redshift radio galaxies. The observations suggest the following scenario: A quasar nucleus, hidden along our line of sight, is responsible for the jets that power the radio source as well as for the illumination of material in the immediate environment of the radio galaxy that falls within a biconical region. This illumination is made evident to us by scattering by dust and by emission from the large Ly- α nebula that is aligned with the radio axis (McCarthy *et al.* 1990). The bulge is apparently still in the process of being assembled from small (~ 1.5 kpc = $0''.2$), merging, dusty objects (*e.g.*, Baron & White 1987; Pascarella *et al.* 1996), which, at the depth of our current images, are visible only by scattered light, when they happen to fall within the illumination cone of the central source. Deeper high-resolution imaging might pick up intrinsic emission from similar objects in regions not illuminated by the quasar.

We are grateful for the support of the University of Hawaii Adaptive Optics group: François Roddier, Claude Roddier, Buzz Graves, Malcolm Northcott, and Laird Close, without which these observations would not have been possible. We also thank Laird Close and Claude Roddier for helpful discussions and for information on the image scale,

Dave Monet and Dave Tholen for discussions on astrometric matters, and Rob Whitely for obtaining short exposures on the 3C 294 field. This research was supported in part by NSF grant AST 95-29078.

REFERENCES

- Baron, E., & White, S.D.M. 1987, *ApJ*, 322, 585
- Chambers, K.C. 1999*a*, in “After the Dark Ages: When Galaxies Were Young,” ed. S.S. Holt & E.P. Smith (AIP Press, College Park, MD), in press
- Chambers, K.C. 1999*b*, *ApJ*, submitted
- Fosbury, R.A.E., Vernet, J., Villar-Martin, M., Cohen, M.H., Cimatti, A., di Serego Alighieri, S., & McCarthy, P.J. 1999, in “NICMOS and the VLT,” ESO Conference and Workshop Proceedings No. 55, ed. W. Freudling & R. Hook (ESO, Garching), p. 187
- Graves, J., Northcott, M., Roddier, F., Roddier, C., & Close, L. 1998, *Proc. SPIE*, 3353, 34
- Hodapp, K.-W., *et al.* 1996, *New Astr.*, 1, 177
- Kristian, J., Sandage, A., & Katem, B. 1974, *ApJ*, 191, 43.
- McCarthy, P.J., Spinrad, H., van Breugel, W., Liebert, J., Dickinson, M., Djorgovski, S., & Eisenhardt, P. 1990, *ApJ*, 365, 487
- Pascarelle, S.M., Windhorst, R.A., Keel, W.C., & Odewahn, S.C. 1996, *Nature*, 383, 45
- Pentericci, L., Röttgering, H.J.A., Miley, G.K., McCarthy, P., Spinrad, H., van Breugel, W.J.M., & Macchetto, F. 1999, *A&A*, 341, 329
- Pentericci, L., Röttgering, H.J.A., Miley, G.K., Spinrad, H., McCarthy, P.J., van Breugel, W.J.M., & Macchetto, F. 1998, *ApJ*, 504, 139
- Riley, J. M., Longair, M. S., & Gunn, J. E. 1980, *MNRAS*, 192, 233
- Roddier, F., Northcott, M., & Graves, J.E. 1991, *PASP*, 103, 131
- Stockton, A., Canalizo, G., & Close, L. 1998, *ApJ*, 500, L121
- van Breugel, W.J.M., Stanford, S.A., Spinrad, H., Stern, D., & Graham, J.R. 1998, *ApJ*, 502, 614
- Veron, P. 1966, *ApJ*, 144, 861
- Wainscoat, R.J., & Cowie, L.L. 1992, *AJ*, 103, 332

Fig. 1.— Image of 3C 294 (centered) in the K' filter obtained with the University of Hawaii AO system Hokupa'a at the CFHT. The wings and diffraction spike from the AO guide star (placed outside the field of the detector) are visible to the right. The compact bright spot just below and to the left of 3C 294 is a 10^{-4} -intensity ghost image of the guide star produced by the optics of the AO system. The FWHM of the image is $\sim 0''.15$. The upper right inset shows 3C 294 at lower contrast, and the upper-left inset shows a short exposure on the AO guide star, which is found to be double, with a separation of $0''.13$ and an intensity ratio at K' of 1.5:1. The large lower-left inset reproduces the region around and to the east of 3C 294, with estimates of the position of the radio nucleus shown. The large cross indicates the position and 2σ internal error range of the radio core based on a position for the AO guide star from an astrometric fit to 12 USNO-A2.0 stars. The three “ $\times$ ” points show the position of the radio core from positions for the AO guide star given by Kristian *et al.* (1974), Veron (1966), and Riley *et al.* (1980) (left to right, respectively). Assuming that the radio core actually lies near the southern apex of the optical structure, the short arrows in the main panel are at the locations of the inner knots of the radio jets and point towards the directions of the hot spots, as seen on the 6 cm VLA map of McCarthy *et al.* (1990).

This figure "fig1.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9905154v1>